


R and R<sup>1</sup> are independently selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>2</sub>-C<sub>20</sub> alkynyl, C<sub>2</sub>-C<sub>20</sub> alkoxy carbonyl, aryl, C<sub>1</sub>-C<sub>20</sub> carboxylate, C<sub>1</sub>-C<sub>20</sub> alkoxy, C<sub>2</sub>-C<sub>20</sub> alkenyloxy, C<sub>2</sub>-C<sub>20</sub> alkynyloxy, or aryloxy, each R and R<sup>1</sup> optionally being substituted with C<sub>1</sub>-C<sub>5</sub> alkyl, halogen, C<sub>1</sub>-C<sub>6</sub> alkoxy, or with a phenyl group substituted with halogen, C<sub>1</sub>-C<sub>5</sub> alkyl or C<sub>1</sub>-C<sub>5</sub> alkoxy;

X and X<sup>1</sup> are independently selected from the group consisting of anionic ligands;

B1 and

L and L<sup>1</sup> are selected from the group consisting of nucleophilic carbenes, phosphine, sulfonated phosphine, phosphite, phosphinite, phosphonite, ether, amine, amide, sulfoxide, carbonyl, nitrosyl, pyridine and thioether, wherein at least one of L or L<sup>1</sup> is a nucleophilic carbene. —

pub C2 / 08  
23. (Amended) A method of performing ring closing metathesis, said method comprising contacting a diterminal diene with a catalytic complex under conditions appropriate, and for a time sufficient to produce a cyclic alkene, wherein the catalytic complex has the formula:



wherein

B2 C<sup>1</sup>, C<sup>2</sup> and C<sup>3</sup> are sp<sup>2</sup>-hybridized carbons, wherein either or both of C<sup>1</sup> and C<sup>2</sup> are optionally absent;

M is selected from the group consisting of Os and Ru;

R and R<sup>1</sup> are independently selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>20</sub> alkyl, C<sub>2</sub>-C<sub>20</sub> alkenyl, C<sub>2</sub>-C<sub>20</sub> alkynyl, C<sub>2</sub>-C<sub>20</sub> alkoxy carbonyl, aryl, C<sub>1</sub>-C<sub>20</sub> carboxylate, C<sub>1</sub>-C<sub>20</sub> alkoxy, C<sub>2</sub>-C<sub>20</sub> alkenyloxy, C<sub>2</sub>-C<sub>20</sub> alkynyloxy, or aryloxy, each R and R<sup>1</sup> optionally being substituted with C<sub>1</sub>-C<sub>5</sub> alkyl, halogen, C<sub>1</sub>-C<sub>6</sub> alkoxy, or with a phenyl group substituted with halogen, C<sub>1</sub>-C<sub>5</sub> alkyl or C<sub>1</sub>-C<sub>5</sub> alkoxy;

X is an anionic ligand; and

L is a nucleophilic carbene; and

Ar is an aryl substituent, bonded to M by an η<sup>6</sup> bond. —

Please add claims 27-38.

pub C9  
-- 27. (New) The catalytic complex of claim 9, wherein at least one of the anionic ligands X and X' are independently selected from the group consisting of halide, carboxylate, alkoxy, aryloxy, and alkyl sulfonate.--

-- 28. (New) The catalytic complex of claim 27, wherein at least one of the anionic ligands is chloride.--

-- 29. (New) The catalytic complex of claim 9, wherein the nucleophilic carbene ligand comprises a carbene carbon further bonded to two heteroatoms having electronegativity greater than that of carbon, wherein the heteroatoms are independently selected from the group consisting of nitrogen, oxygen, and sulfur.--

pub D9  
B3  
-- 30. (New) The catalytic complex of claim 29, wherein the nucleophilic carbene ligand comprises a saturated or unsaturated 1,3-diheteroaromatic cyclic compound.--

-- 31. (New) The catalytic complex of claim 9, wherein the complex is linked to a solid support by means of a link between at least one of said anionic ligands and said solid support.--

pub C10  
-- 32. (New) The catalytic complex of claim 9, wherein the complex is linked to a solid support by means of a link between at least one of said nucleophilic carbenes and said solid support.--

-- 33. (New) The catalytic complex of claim 14, wherein the anionic ligand X is selected from the group consisting of halide, carboxylate, alkoxy, aryloxy, and alkyl sulfonate.--

-- 34. (New) The catalytic complex of claim 33, wherein the anionic ligand is chloride.--

-- 35. (New) The catalytic complex of claim 14, wherein the nucleophilic carbene ligand comprises a carbene carbon further bonded to two heteroatoms having electronegativity greater than that of carbon, wherein the heteroatoms are independently selected from the group consisting of nitrogen, oxygen, and sulfur.--

pub D9  
-- 36. (New) The catalytic complex of claim 35, wherein the nucleophilic carbene ligand comprises a saturated or unsaturated 1,3-diheteroaromatic cyclic compound.--